

# Guide to iron supplementation in pregnancy

This publication can be used by GPs, Midwives and Maternity Professionals

Date of publication: September 2021

FOR HEALTH PROFESSIONALS ONLY



ISTOCKPHOTO.COM/MIHALILOVANOVIC

In association with:



**Complete your CPD using one of the following steps...**

**Visit;**  
[healthprofessionalacademy.co.uk/  
iron-supplements](https://healthprofessionalacademy.co.uk/iron-supplements)



**Scan this QR code to take  
this CPD**



# Iron deficiency is common in pregnancy, with risks to mum and baby

The demand for iron increases threefold during pregnancy to meet the demands of the growing fetus and placenta, as well as maternal adaptations to pregnancy.<sup>1</sup> It is estimated that 30-40% of pregnant women in industrialised countries are iron deficient<sup>1</sup> and almost one in four women in the UK have iron deficiency anaemia.<sup>2</sup>

Iron deficiency during pregnancy is associated with several complications including increased risk of preterm birth, lower birth weight and perinatal mortality.<sup>3,4</sup> It is also linked to maternal postpartum fatigue and depressive symptoms, and infant iron deficiency in the first three months of life.<sup>2,3</sup> Treating iron deficiency therefore offers considerable benefits to both mother and baby.

## Iron is recycled between functional and storage ‘pools’

Around a quarter of the body’s iron is stored as ferritin, mainly in the liver, ready to be mobilised if the body requires additional iron. The remaining iron is mostly ‘functional’ iron, found predominantly in haemoglobin (Hb) in circulating red blood cells but also in myoglobin in skeletal muscle. A small fraction (< 1%) circulates in the blood bound to transferrin; this is known as ‘transport’ iron.<sup>3-5</sup>

## Iron stores are depleted before anaemia develops

Iron deficiency can be viewed as a spectrum from iron depletion through to iron deficiency anaemia dependent on the status of stored iron in the body (Figure 1).




Figure 1: The spectrum of iron deficiency

**ABOUT THE HEALTH PROFESSIONAL ACADEMY**  
The Health Professional Academy (HPA) is a free online learning channel from Lifecycle Marketing, the publishers of *Emma's Diary* and parenting website, *Families*. We've been a trusted partner in healthcare education for more than 28 years, working alongside professional health bodies and Royal Colleges.

All our online CPD modules are developed in partnership with a team of leading academics, so you can be sure the module you've chosen is credible, referenced and up to date. All information by HPA is in line with guidance from the NHS, NICE and Royal Colleges, as well as Unicef's Baby Friendly Initiative.

**HEALTH PROFESSIONAL ACADEMY TEAM**  
Healthcare Director: Sunil Singh  
Reviewed by Amanda Burleigh RGN, RM, BSc (hons).

**CONTACT US**  
Should you have any questions relating to the material within this module or general comments about HPA, please do not hesitate to contact us at:

✉ [info@healthprofessionalacademy.co.uk](mailto:info@healthprofessionalacademy.co.uk)  
☎ 01628 535499

**ACKNOWLEDGEMENT**  
HPA materials are produced for training purposes of HEALTH PROFESSIONALS only. No endorsement of the products in the module or across any of our channels is implied or intended by HPA, its authors or reviewers. Materials are not to be used or shared beyond their intended use without prior consent from the Health Professional Academy.

Any adverse event arising from the use of a product should be reported directly to the product manufacturer. Adverse events arising from medicines or medical devices should also be reported through the Yellow Card Scheme. Reporting forms and information can be found at <https://yellowcard.mhra.gov.uk>

© Health Professional Academy 2021

Lifecycle Marketing (M&B) Ltd, Remenham House, 3 Regatta Place, Marlow Road, Bourne End, Buckinghamshire, SL8 5TD.  
Registered in England No: 2556692

In iron depletion, levels of stored iron are reduced, while levels of circulating iron in Hb may not be affected.<sup>3</sup> In iron-deficiency anaemia, red blood cell production is diminished due to a shortage of both stored and circulating iron.<sup>3</sup>

Stored iron is therefore depleted before Hb decreases. Since iron is essential to all cells, symptoms of iron deficiency – such as fatigue, irritability, poor concentration and hair loss – may occur even in the absence of anaemia.<sup>4</sup>

Many women, especially those who experience heavy menstrual bleeding, already have low iron reserves at the time of conception, putting them at even greater risk of iron deficiency during pregnancy. Indeed, it is estimated that 10-20% of menstruating women are iron deficient.<sup>4</sup>

### Current UK guidance on iron deficiency in pregnancy

The British Committee for Standards in Haematology (BCSH) published guidelines on the management of iron deficiency in pregnancy in 2012.<sup>3</sup> These are reflected in the latest NICE guidance on iron deficiency anaemia.<sup>2</sup>

The BCSH guidelines recommend a full blood count at booking and at 28 weeks. Normal Hb levels vary during pregnancy, therefore anaemia is defined as:<sup>3</sup>

- Hb <110g/L in the first trimester
- Hb <105g/L in the second and third trimesters
- Hb <100g/L in the post-partum period.

Women who are not anaemic but thought to be at increased risk of iron deficiency (see Box 1) should have their serum ferritin levels checked early in pregnancy. Serum ferritin levels progressively fall by 32 weeks to around 50% of pre-pregnancy levels due to haemodilution as plasma volume increases as well as mobilisation of iron.<sup>2-4</sup>

Even with this gradual decline in serum ferritin, levels <15ug/L indicate iron deficiency in all stages of pregnancy. Levels <30ug/L indicate early iron depletion, which will continue to fall during pregnancy unless treated.<sup>2,3</sup>

### Dietary changes alone are insufficient to correct iron deficiency in pregnancy

Once a woman becomes iron deficient during pregnancy, dietary sources of iron will be insufficient to replenish levels and oral iron supplementation is recommended.<sup>2,3,7</sup>

NICE guidelines recommend that pregnant women with iron deficiency anaemia should be prescribed 200mg oral ferrous sulphate tablets two or three times a day. Treatment should continue for three months after the deficiency has been corrected to ensure iron stores are replenished.<sup>2</sup>

Hb levels (full blood count) should be rechecked after two to four weeks to assess response to treatment.<sup>2</sup>

Women with serum ferritin levels below 30ug/L should be offered iron supplements, as this indicates early iron depletion, which will continue to fall unless treated.<sup>2</sup>

### Box 1: Women at higher risk of iron deficiency during pregnancy<sup>2,8</sup>

- Low iron stores before pregnancy
- Previous history of anaemia
- Pre-existing blood condition (e.g. sickle cell disease, thalassaemia)
- Age < 20 years
- Dietary deficiency: gut disorders affecting iron absorption (e.g. coeliac disease, inflammatory bowel disease, previous gut surgery); eating disorders; vegetarian/vegan
- Short spaces between pregnancies (< 18 months)
- Multiple pregnancy

## Treating iron deficiency can be challenging

There are many non-haem iron supplements available over the counter. Ferrous sulphate is considered the 'gold standard' and is the most commonly prescribed form in the UK.<sup>4</sup> Oral iron supplements, however, can have limitations: only a small percentage of the iron is absorbed (around 10-15%)<sup>10</sup> and side effects are common, which may in turn lead to poor compliance.

Bioavailability among supplements varies, and it is important to distinguish between the amount of iron compound and the equivalent in terms of elemental iron. For example, hydrated ferrous sulphate contains 20% iron by weight, therefore a 300mg tablet contains 60mg of elemental iron.<sup>4</sup>

### Iron is absorbed via a carrier mechanism, DMT-1

Although some nutrients can enter the body by passing between the gut mucosal cells, iron uptake and transfer depends on specific cellular carrier mechanisms. One of these, called the divalent metal transporter 1 (DMT-1), plays an important role in the uptake of non-haem iron.<sup>4</sup>

The absorption of most dietary iron occurs in the duodenum and upper jejunum, where DMT-1 is found on the surface of the absorptive cells (enterocytes) that line the small intestine (Figure 2).<sup>11</sup> The acidic pH in the duodenum is critical to the absorption of iron.<sup>11</sup>

Various factors can affect the absorption of iron supplements (see Box 2).




Figure 2: Absorption of iron

### Box 2. Factors affecting the absorption of iron supplements

- **Solubility of the iron:** ferrous iron is more easily absorbed than ferric iron.<sup>9,12</sup>
- **Timing of the dose:** taking iron supplements with or after food may decrease absorption by 40-66%.<sup>2,11</sup>
- **Dietary factors:** iron absorption is inhibited by polyphenols (found in various plant-based food and beverages such as tea, coffee and chocolate), phytates (also found in plant-based diets), calcium, legumes, cereals and some animal and plant proteins;<sup>5,11</sup> conversely, iron absorption is increased in the presence of vitamin C.<sup>5,11</sup>
- **Other medication:** drugs that alter gastric acid production (e.g. proton pump inhibitors) can also substantially reduce iron absorption.<sup>4,11</sup>



ISTOCKPHOTO.COM/KALI9

### **Side effects are common and affect compliance**

Iron has a highly reactive chemical nature, which allows it to fulfil its essential functions within the body. However, this also gives it the ability to cause oxidative stress and cell damage within the gastrointestinal tract.<sup>13</sup>

Although iron absorption occurs mainly in the small intestine, traditional iron supplements tend to dissolve in the stomach or, in the case of delayed release products, the lower intestine where they can cause damage not only to the gut lining but also to the gut microbiome.<sup>9,14</sup> This can lead to inflammation and the potential for side effects including constipation, diarrhoea, heartburn, epigastric pain, faecal impaction, gastrointestinal irritation and nausea.<sup>2,13</sup>

Estimates of the incidence of gastrointestinal side effects with iron supplementation range from 30-47%.<sup>15</sup>

In pregnancy, the side effects of iron supplementation may be compounded by those of the growing fetus, such as nausea, heartburn and constipation.<sup>9,15</sup> Indeed, one study confirmed that pregnant women were more likely to report side effects when taking daily iron supplements than controls (25.3% versus 9.9%), particularly at doses of 60mg or more of elemental iron.<sup>16</sup>

Adverse effects are a common cause of non-compliance with treatment: up to 50% of patients are thought to discontinue iron supplements because of adverse effects.<sup>9</sup>

# Active Iron offers a unique formula to increase absorption and reduce side effects

Active Iron's ground-breaking formula delivers iron to the body's normal site of absorption in the duodenum, the DMT-1. This not only increases the amount of iron absorbed, it also helps protect the gut lining from iron-related oxidative stress and damage, resulting in fewer side effects.<sup>14</sup>

## Better iron absorption than ferrous sulphate

Clinical evaluation of Active Iron showed significantly greater absorption of iron when compared with the gold standard of ferrous sulphate alone.<sup>14</sup> In this controlled, pharmacokinetic study of 21 healthy participants who were mostly women, iron absorption with Active Iron was double that of ferrous sulphate alone (Figure 3).<sup>14</sup>


Because iron absorption is dependent on a

person's degree of iron depletion, further analysis was carried out. Multi-variable analysis showed that the greater absorption of Active Iron was independent of baseline serum iron, ferritin, transferrin and haemoglobin levels.<sup>14</sup>


## Less gut irritation resulting in improved compliance and good efficacy

When iron is not fully absorbed in the gut, it stays in its ferric state ( $\text{Fe}^{3+}$ ). This can lead to gastrointestinal side effects such as constipation and nausea. Active Iron is different. Active Iron is delivered as ferrous iron ( $\text{Fe}^{2+}$ ), providing better bioavailability and reduced side effects as a result.

Further clinical evaluation found that Active Iron was six times less likely to cause gut irritation compared to previous oral iron, resulting in four times better compliance in taking the product.<sup>17</sup> In this randomised, double-blind study of 59 women with a history of intolerance to oral iron, Active Iron showed a 94% increase in iron stores (Figure 4), with a significant improvement in resulting energy levels.<sup>17</sup>



**Figure 3:** Active Iron achieves twice the absorption of ferrous sulphate alone<sup>14</sup> ( $P=0.01$ ).




**Figure 4:** Active Iron showed a 94% increase in median Ferritin levels (from 8.00 to 15.5  $\mu\text{g/L}$ ) in the overall cohort, up within 6 weeks and sustained at 12 weeks<sup>17</sup> ( $P=0.0002$ ).

## Active Iron Pregnancy Plus offers a complete nutritional solution for mum & baby

Available over-the-counter, Active Iron Pregnancy Plus is a specially developed, combination product containing 3 components:

- Non-constipating Active Iron capsule. Clinically proven to have 2X better absorption versus standard ferrous sulfate. Active Iron's formula increases iron and helps to reduce gut irritation from iron.<sup>14, 17</sup>
- 100% sustainably sourced omega-3 DHA fish oil capsule, with AlaskOmega® omega-3 concentrates.
- A specially designed multivitamin tablet with Vitamin D and Active Folic, a source of folic acid in an active, naturally occurring form that delivers greater absorption compared to standard folic acid.

Pregnant and postpartum women should take one Active Iron capsule, one multivitamin with Active Folic tablet and one omega-3 DHA capsule daily.



Food supplements are not a substitute for a varied diet and healthy lifestyle.

## i References

1. UN Children's Fund, UN University, World Health Organization. Iron deficiency anaemia: assessment, prevention and control. A guide for programme managers, 2001.
2. NICE. Anaemia – iron deficiency. September 2018.
3. Pavord S et al. UK guidelines for the management of iron deficiency in pregnancy. *British Journal of Haematology* 2012; 156(5): 588-600.
4. Scientific Advisory Committee on Nutrition. Iron and health. 2010. [www.sacn.gov.uk](http://www.sacn.gov.uk) Accessed March 2019.
5. Abbaspour N et al. Review on iron and its importance for health. *J Res Med Sci* 2014; 19 (2): 164-174.
6. Suppi ET. Iron deficiency without anaemia – a clinical challenge. *Clinical Case Reports* 2018; 6 (6): 1082-1086.
7. Goddard AF et al. Guidelines for the management of iron deficiency anaemia. *Gut* 2011; 60: 1309-1316.
8. NHS. Iron deficiency anaemia in pregnancy: information for patients. January 2017.
9. Tolken Z et al. Ferrous sulphate supplementation causes significant gastrointestinal side effects in adults: a systematic review and meta-analysis. *PLoS ONE* 2015; 10 (2): e0117383 doi: 10.1371/journal.pone.0117383
10. Santiago P. Ferrous versus ferric oral iron formulations for the treatment of iron deficiency: a clinical overview. *The Scientific World Journal* 2012, Article ID 846824. Doi:10.1100/2012/846824
11. Ems T, Huecker MR. Biochemistry, Iron Absorption. [Updated 2018 Oct 27]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2019 Available from: <https://www.ncbi.nlm.nih.gov/books/NBK448204/>
12. Institute of Medicine (US) Committee on Nutritional Status During Pregnancy and Lactation. Nutrition during pregnancy. Washington DC: National Academic Press 1990.
13. Brannon P, Taylor CL. Iron supplementation during pregnancy and infancy: uncertainties and implications for research and policy. *Nutrients* 2017; 9. Doi: 10.3390/nu9121327.
14. Wang J et al. Novel iron-whey protein microspheres protect gut epithelial cells from iron-related oxidative stress and damage and improve iron absorption in fasting adults. *Acta Haematol* 2017; 138: 223-232. Doi: 10.1159/000480632
15. Cancelo-Hidalgo MJ et al. Tolerability of different oral iron supplements: a systematic review. *Curr Med Res Opin* 2013; 29 (4): 291-303. Doi: 10.1185/03007995.2012.761599.
16. Pena-Rosas JP et al. Intermittent oral iron supplementation during pregnancy. *Cochrane Database Syst Rev* 2012, 7:CD009997. Doi:10.1002/14651858.
17. Ledwidge et al. Low iron, iron deficiency and anaemia in adult women with self-reported oral iron gastrointestinal intolerance and management with an iron-whey-protein formulation. Data on file.



# ACTIVEIRON®

## PREGNANCY PLUS

Complete nutritional solution  
for mum and baby



**NON-CONSTIPATING  
ACTIVE IRON 25MG**

**SUSTAINABLY SOURCED  
OMEGA-3 DHA**

**COMPLETE MULTIVITAMIN  
WITH ACTIVE FOLIC**

REQUEST A FREE SAMPLE

[info.activeiron.com/pregnancy-sample-request](https://info.activeiron.com/pregnancy-sample-request)